

NEUTRINO PROPERTIES AND SUSY WITHOUT R -PARITY *

Ralf Hempfling

*Max-Planck-Institut für Physik, Werner-Heisenberg-Institut,
Föhringer Ring 6, 80805 Munich, Germany*

and

Univ. of California at Davis, Dept. of Physics, Davis, CA 95616

E-mail: hempf@bethe.ucdvis.edu

ABSTRACT

In supersymmetric models without R -parity neutrinos naturally become massive and mix with each other. We explore the predictions of a very restricted model with only three free parameters and find that this model naturally yields masses and mixing angles compatible with experimental results from solar and atmospheric neutrino experiments. Furthermore, there is a tiny region in parameter space where the solution to the solar neutrino puzzle is compatible with either the LSND result or the existence of significant hot dark matter neutrinos.

* to be published in the Preceedings of the VIIIth Rencontres de Blois: "Neutrinos, Dark Matter and the Universe", Blois, France (June 8-12, 1996)

In the Standard Model of elementary particles (SM) both lepton number (L) and baryon number (B) are conserved due to an accidental symmetry, *i.e.* there is no renormalizable, gauge-invariant term that would break the symmetry. In the minimal supersymmetric extension of the SM (MSSM) the situation is different. Due to a the variety of scalar partners the MSSM allows for a host of new interactions many of which violate B or L .

Since neither B nor L violation has been observed in present collider experiments these couplings are constrained from above. More constraints arise from neutrino physics or cosmology. Thus, all lepton and baryon number violating interaction are often eliminated by imposing a discrete, multiplicative symmetry called R -parity,¹ $R_p \equiv (-1)^{2S+3B+L}$, where S is the spin. One very attractive feature of R_p conserving models is that the lightest supersymmetric particle (LSP) is stable and a good cold dark matter candidate.²

However, while the existence of a dark matter candidate is a very desirable prediction, it does not prove R_p conservation and one should consider more general models. Here, we will investigate the scenario where R_p is broken explicitly via the terms³ $W = \mu_i L_i H$, where H is the Higgs coupling to up-type fermions and L_i ($i = 1, 2, 3$) are the left-handed lepton doublets. Clearly, these Higgs-lepton mixing terms violate lepton-number. As a result, majorana masses will be generated for one neutrino at tree-level and for the remaining two neutrinos at the one-loop level. These masses were calculated in the frame-work of minimal supergravity in ref. 4 and the numerical results will be briefly summarized here.

There are three R_p violating parameters which can be used to fix 1) the tree-level neutrino mass, 2) the μ - τ mixing angle and 3) the e - μ mixing angle. The question of whether *e.g.* the solar⁵ and the atmospheric⁶ neutrino puzzle can be solved simultaneously depends on the prediction of $m_{\nu_\mu}^2 - m_{\nu_e}^2$. In fig. 1 we have scanned the entire SUSY parameter space consisting of the Higgsino (gaugino) mass parameter, μ ($m_{1/2}$), the trilinear scalar interaction parameter A_0 , and the ratio of Higgs VEVs, $\tan\beta$. The universal scalar mass parameter m_0 is fixed by minimizing the potential. Plotted is the number of models yielding a particular prediction for $m_{\nu_\mu}^2 - m_{\nu_e}^2$ for (a) $\sin^2 2\theta_{e\nu_\mu} = 0.008$ and (b) $\sin^2 2\theta_{e\nu_\mu} = 1$. We fix $m_{\nu_\tau} = 0.1$ eV and $\sin^2 2\theta_{\mu\nu\tau} = 1$ in order to solve the atmospheric neutrino problem. We see that both long wavelength oscillation (LWO)⁷ ($m_{\nu_\mu}^2 - m_{\nu_e}^2 = 10^{-10}$ eV²) and MSW effect⁸ ($m_{\nu_\mu}^2 - m_{\nu_e}^2 = 10^{-5}$ eV²) can be accommodated. In fig. 1(c) we solve the solar neutrino problem via e - τ oscillations and we fix $\sin^2 2\theta_{e\nu_\mu} = 0.004$ in order to accommodate the LSND result.⁹ We see that most models are already ruled out by collider constraints and even more by dark matter (DM) constraints. However, a very small (but non-zero) number of models yields a prediction compatible with the LSND result (the dotted line is lower limit of LSND).

Acknowledgements This work was supported in parts by the DOE under Grants No. DE-FG03-91-ER40674 and by the Davis Institute for High Energy Physics.

References

1. N. Sakai and T. Yanagida, *Nucl. Phys.* **B197**, 533 (1982).
2. J. Ellis *et al.*, *Nucl. Phys.* **B238**, 453 (1984).
3. L.J. Hall and M. Suzuki, *Nucl. Phys.* **B231**, 419 (1984).
4. R. Hempfling, MPI-PhT/95-59, hep-ph/9511288, *Nucl. Phys.* **B**, to appear.
5. P. Anselmann *et al.*, *Phys. Lett.* **B327**, 234 (1994).
6. Y. Fukuda *et al.*, *Phys. Lett.* **B335**, 237 (1994).




Fig. 1. Histogram of the number of models that yield a particular prediction for $m_{\nu_\mu}^2 - m_{\nu_e}^2$ assuming (a) small angle and (b) large angle solution to solar neutrino problem. In (c) we solve the solar neutrino problem via small angle e - τ oscillations and check whether this is compatible with the LSND result.

7. V. Gribov and B. Pontecorvo, *Phys. Lett.* **B28**, 493 (1969); V. Barger, R.J.N. Phillips and K. Whisnant, *Phys. Rev.* **D24**, 538 (1981); *Phys. Rev. Lett.* **69**, 3135 (1992).
8. L. Wolfenstein, *Phys. Rev.* **D17**, 2369 (1978); **20**, 2634 (1979); S.P. Mickheyev and A. Yu Smirnov, *Yad. Fiz.* **42**, 1441 (1985) [*Sov. J. Nucl. Phys.* **42**, 913 (1986)].
9. C. Athanassopoulos *et al.* , *Phys. Rev.* **D75**, 2650 (1995); LA-UR-96-1326.